Clinic Sofia **Prenatal Information Packet**

Thank you for choosing Clinic Sofia OB/GYN, PA for your obstetrical care. We are happy to be sharing this exciting time with you! We have taken the time to put together this packet for you, so please read through carefully and thoroughly, as it includes information on what to expect throughout pregnancy, appointments, screenings, billing information, delivery, registration with the hospital, and postpartum depression. Hopefully, this packet of information will be a helpful reference for you in the upcoming months. If you have any additional questions, please call the main line at (952) 922-7600.

The clinic hours are Monday 8:30 am to 5:00 pm, Tuesday through Thursday 8:00 am to 5:00 pm, and Friday 8:00 am to 12:00 pm. After office hours, emergency calls will be handled by the physician on call who can be reached by calling the main line at (952) 922-7600.

www.clinicsofia.com

Clinic Sofia – Edina

6545 France Avenue South, 490 Edina, MN 55435 952-922-7600 Phone 952-345-4448 Fax Clinic Sofia – Maple Grove 15715 Grove Circle North Maple Grove, MN 55369 952-922-7600 Phone 952-345-4448 Fax

Jewelia D Wagner, MD Amy L Hammers, MD David F Clay, MD Pamela A Jordi, MD Staci Biegner, MD Jennifer Knapp, MSN, WHCNP Sarah Iverson, PA-C Elise Pesch, DNP, FNP-BC Sarah Talbott, PA-C Welcome to your pregnancy confirmation visit and congratulations on your pregnancy! This packet is a general guideline of what to expect in your pregnancy and how we at Clinic Sofia will support you through your journey. Much of this information – and more – is available on our website which is listed on the first page. We look forward to sharing this experience with you and working towards a healthy and safe delivery.

PROVIDERS, ON-CALL, HOSPITALS:

We currently have 5 doctors and 4 advanced practice providers. We do have a call schedule and cover each other when on call. We do not share call with any other groups or providers, so you will have a Clinic Sofia doctor at your deliver. While we are excited to meet your new baby and will attempt to be at each of our patients' deliveries, we cannot guarantee this to be the case. We plan for you to meet all the providers, so you will always have a familiar face at the time of delivery. We have privileges at two hospitals, **Maple Grove Hospital** and **Fairview Southdale Hospital**.

PRENATAL OFFICE VISITS:

Below is an example of the prenatal schedule for uncomplicated pregnancies. Earlier in your pregnancy, your visits may take more time while later visits may be more concise. If your pregnancy has complications, you will be seen more frequently.

WEEK 5-13 (first trimester)

You can expect to be seen roughly every 3-4 weeks in the first trimester. We will assess your vital signs, listen to fetal heart tones via a doppler device, discuss symptoms and expectations, and answer specific questions you may have. A few details to prepare you:

 1^{st} visit: Blood will be drawn at this visit for your blood type, antibody levels, hemoglobin, and sexually transmitted infections. If you are > 7 weeks gestation, an ultrasound will be performed to confirm gestational age, heartbeat, number, and location of the pregnancy. If you are < 7 weeks, a separate visit will be scheduled with an ultrasound.

2nd visit: A breast and pelvic exam will be completed. If you are due for a pap smear, we can safely collect at this visit. A swab for gonorrhea and chlamydia is also routinely collected. A blood draw can be collected for genetic screening if you have chosen to do so.

WEEKS 14-27 (second trimester)

You can expect to be seen roughly every 4 weeks in the second trimester. We will assess your vital signs, listen to fetal heart tones via a doppler device, discuss symptoms and expectations, and answer specific questions you may have. A few details to prepare you:

~16-week visit: If you choose to screen for Spina Bifida, your blood will be drawn for the AFP (Alpha Fetal Protein) lab.

~20-week visit: A thorough ultrasound of fetal anatomy is performed. You will review the results with your physician after the ultrasound. If the pregnancy is low risk, no other ultrasounds may be indicated in your pregnancy.

WEEKS 28-40 (third trimester):

You can expect to be seen roughly every 2 weeks initially, and then you will be seen weekly starting at 36 weeks. We will assess your vital signs, listen to fetal heart tones via a doppler device, measure fundal height, discuss symptoms and expectations, and answer specific questions you may have. A few details to prepare you:

~28-week visit: Expect to be here for 1 hour as we will complete your diabetes screening, additional testing may be warranted based on the result. We routinely check your blood count and a syphilis test with a blood draw. It is recommended to receive the TDAP vaccine during this visit.

~36-week visit: We will collect a vaginal swab to assess for Group B strep. We may recommend cervical exam to verify fetal position.

**Full term: 37-40-week visits: If you desire, your provider can perform a cervical check to assess dilation.

Additional visits and/or ultrasounds may be added depending on your specific pregnancy needs.

WHEN YOU GO INTO LABOR, CALL THE CLINIC DAY OR NIGHT: 952-922-7600

NORMAL CHANGES IN PREGNANCY:

NAUSEA AND VOMITING IN PREGNANCY:

To prevent nausea, try the following suggestions until you find one that works.

- Eat a piece of bread or a few crackers before you get out of bed in the morning or when you feel nauseated.
- Get out of bed slowly. Avoid sudden movements.
- Have some yogurt, cottage cheese, juice or milk before you go to bed or before you get up. Try one of these if you have to get up during the night. Have a protein snack at bedtime and in the morning.
- Eat several small meals during the day so your stomach doesn't remain empty for very long.
- Eat high protein foods (ex. eggs, cheese, nuts, meat, etc) as well as fruits and fruit juices. These foods help prevent low levels of sugar in your blood which can also cause nausea.
- Drink soups and other liquids between meals instead of with meals. Wait about 20" after you have eaten solids to drink the liquids.
- Avoid greasy or fried foods. They are harder to digest.
- Avoid spicy, heavily seasoned foods.

To remedy nausea/vomiting, try these suggestions:

- "Sea-bands" placed on the wrist have been shown to be effective.
- Sip soda water when you begin to feel nauseated.
- Take deep breaths.
- Drink spearmint, raspberry, or peppermint tea. Ginger tea/lozenges may be helpful as well.
- Unisom and Vitamin B6 as directed on page 7.

CONSTIPATION:

Constipation is a very common condition affecting pregnant women. Since this condition is so common, we have printed this sheet to help you achieve natural control of this problem. The digestive tract extracts the beneficial elements from food and then must eliminate the remainder as waste. The whole process takes 1-3 days, the time varying from one individual to another. Hormonal changes may slow this process as may the addition of iron in the prenatal vitamin exacerbates the condition.

The colon, also known as the large intestine, is the last 5 feet of the intestinal tract whose job is to absorb excess water from the fecal waste. This waste is propelled through the entire digestive tract by muscular contractions. If the contractions are too fast, bowel movements will be liquid and if they are too slow, bowel movements will be hard. To prevent constipation, the transit time and moisture level must be correct. Diet is very important to achieving balance. High residue foods, especially those with high fiber content, help provide the bulk to push waste through and to hold on to the necessary moisture. Foods high in fiber include:

- Bran, whole grains, cereals
- Raw fruit (apples, melon, peaches, etc.)
- Cooked or dried fruit
- Raw vegetables (broccoli, cabbage, celery, etc.) or cooked vegetables

Some Helpful Habits for Normal Bowel Movements

- Eat at regular hours and chew food thoroughly
- Drink plenty of fluid
- Exercise daily
- Answer the urge to have a bowel movement when it comes

A high fiber diet may be an important part to preventing colon cancer. As you gradually increase the fiber in your diet, foods like bran and cabbage may initially produce an increase in gas and bloating. This tendency will usually subside in a few weeks. You should also try various high fiber foods to see which best agree with you. Another way of adding fiber and bulk to your diet is the use of Metamucil. This product is a natural grain product made from the husk of psyllium seeds. It is also sold under less expensive generic labels. Metamucil is a simplified way of normalizing bowel function that does not require you to radically change your diet. To be effective, however, you must take it daily.

ADDITIONAL NORMAL CHANGES IN PREGNANCY:

Because of your increased blood volume and/or high levels of estrogen and progesterone, the following are additional common complaints that are normal in pregnancy:

- Shortness of breath
- Urinary frequency
- Fatigue
- Feet swelling
- Nasal dryness, stuffiness
- Nose bleeds
- Gums bleeding
- Insomnia
- Extremities falling asleep
- Varicose veins
- Corneal changes eyes dry
 - Don't have an eye exam as the shape of the cornea is changed until 3 months after delivery or 6 weeks after breast feeding is discontinued
- Headaches
- Rib pain
- Dizziness
- Round ligament pain
- Back pain
- Breast pain

POTENTIALLY CONCERNING, ABNORMAL CHANGES IN PREGNANCY:

Complaints that you need to call the office about would be concerns about the previous mentioned things and:

- Persistent headache, especially after 28 weeks
- Blurred or spotty vision of flashes of light
- Sudden swelling of the face and hands
- Inability to keep food or liquids down for two days

If your personal history is significant for risk factors for Pre-Eclampsia, your provider may recommend a baby Aspirin (81 mg) daily from 12 weeks through delivery.

COMMONLY ASKED SAFETY CONCERNS:

Safe during pregnancy:

- Exercise this is encouraged during pregnancy so long as violent bouncing (e.g., trampolines) is avoided
- Fish consumption two to three servings of fish with a high DHA and low mercury content is safe
- Hair dyes although data is limited, systemic absorption of hair dye is minimal and thus presumed safe
- Sexual intercourse provided the patient is not bleeding, doesn't have a placenta previa, and the amniotic membranes are still intact (i.e., water isn't broken), then there is no restriction regarding sexual intercourse
- Travel airline travel is safe though destinations may have infection risks. While there is no exact gestation age at which patients must stop travel, this should be discussed with your doctor prior to travel in 3rd trimester
- Swimming swimming pool usage is safe in pregnancy (though see below for hot tubs)

Decrease during pregnancy:

- Artificial sweeteners data regarding saccharin is conflicting, though low consumption is likely safe
- Caffeine low-to-moderate intake does not appear to be associated with adverse outcomes

Avoid during pregnancy:

- Hot tubs although data is limited, hot tub usage should be avoided in 1st trimester due to spina bifida risk
- Alcohol there is no known threshold between safe and unsafe amount of alcohol in pregnancy and therefore should be avoided
- Smoking, vaping, nicotine women should not smoke cigarettes during pregnancy though if unable to quit entirely, attempt to reduce as much as possible and nicotine replacement (lozenge, gum) is appropriate to help
- Marijuana while there are no known adverse outcomes, long-term neurodevelopmental data is lacking

OB ULTRASOUND:

Occasionally, we will perform an ultrasound earlier if there is a question about the due date or if there is a history of recurrent miscarriage. We will recommend a diagnostic ultrasound for most obstetric patients usually from **18-22** weeks.

The examination consists of placing a special gel on the mother's abdomen and then moving a transducer across it. The returning echoes are electronically converted to a picture on a screen which can then be photographed. Actual fetal movements can be seen. Only <u>ONE</u> guest over the age of 18 (besides the patient) will be allowed into the room. No video, camera, or recording devices are allowed. If recording devices are being used, the ultrasound exam will be discontinued.

Obstetric ultrasound is safe and has become so useful that we feel its routine usage is necessary to provide you with the best prenatal care. Other ultrasounds may be ordered by your provider if deemed medically necessary.

A routine exam usually shows:

- The number of babies
- The fetal age
- The location of the placenta
- Fetal life (the pulsating heart is usually visible)
- Fetal position

Other uses are:

- To help diagnose an early miscarriage
- To detect ovarian cysts and uterine fibroids
- To look for certain congenital malformations and anomalies
- To detect tubal pregnancies
- To investigate abnormal bleeding
- To follow a fetus for growth evaluation

If you have any questions regarding this procedure, please ask your provider.

WEIGHT GAIN DURING PREGNANCY:

Pre-pregnancy weight category	Body Mass Index	Recommended Range of total weight (Ibs)	Recommended range of weight gain in the 2 nd and 3 rd trimesters (Mean Range[lbs/wk])
Underweight	Less than 18.5	28-40	1 (1-1.3)
Normal Weight	18.5-24.9	25-35	1 (0.8-1)
Overweight	25-29.9	15-25	0.6 (0.5-0.7)
Obese (all classes)	30 and greater	11-20	0.5 (0.4-0.6)

**If you have additional questions regarding weight gain during your pregnancy, ask your Provider.

*Reference ACOG and Institute of Medicine: weight gain during pregnancy

INSURANCE:

We provide services for many different health insurance plans; however, it is imperative that you check the requirements of your plan regarding notification of hospitalization and policies regarding length of postpartum stays. Some questions to as yourself and your insurance company before delivery are:

- Do I need pre-approval for my delivery?
- What hospital does my health plan require me to use? (Recall, Clinic Sofia delivers at Fairview Southdale Hospital and Maple Grove Hospital)
- What is my approved length of stay for a vaginal delivery? What about for a Cesarean delivery? Does this time begin at admission to the hospital or at delivery of the baby?
- How do I obtain authorization for a longer stay if either my baby or I have complications? Can I stay if I am doing well but my baby has complications requiring additional days?
- Is circumcision covered? Will the pediatrician be doing the circumcision in the hospital or in their office?
- Are home visits by a nurse covered?
- How is my new baby added to my insurance policy and when does this take effect?
- Are parenting classes covered?
- Will the 20-week ultrasound be covered?
- Will a breast pump be covered?

DISABILITY POLICY FOR PREGNANT PATIENTS:

The great majority of expectant mothers can continue to work until late in pregnancy without any problem. Sometimes, however, the physical changes entailed in pregnancy, or the demands of a woman's job can create workplace difficulties. Please let us know if you have any concerns in this regard. We usually are able to suggest simple steps to deal with the fatigue, morning sickness, or aches and pains that can be common in pregnancy but particularly challenging while you are at work.

If you have more serious symptoms or concerns about potential workplace hazards to you or your baby, we will evaluate the situation and respond accordingly. When medically appropriate, we will recommend that a pregnant patient be placed on disability leave from her job. Such leave is rarely required, though, and in the absence of a serious condition that would endanger the health of the mother or the baby, medical ethics prevent us from making such a recommendation. We will, however, do everything we can to reduce or eliminate pregnancy-related difficulties you may be having at work. This includes contacting your employer, when appropriate, to recommend helpful adjustments or alterations to your duties. Society views pregnancy as a normal condition. Therefore, normal conditions accompanying pregnancy are not covered by disability insurance. Disability insurance coverage applies only to complications of pregnancy. Again, please inform us of any work-related concerns you may have.

SAFE OVER-THE-COUNTER MEDICATIONS IN PREGNANCY AND BREASTFEEDING

PROBLEM	MEDICATION	DOSAGE
Allergy	Allegra Benadryl Claritin Zyrtec	Use as directed on the package
Cold and Congestion	Afrin Nasal Spray Flonase Mucinex	Use as directed on the package, but call the clinic if symptoms do not improve
Constipation	High fiber foods/Metamucil/Psyllium Magnesium supplement Colace (your PNV may contain this) Miralax Milk of Magnesia Senokot	Use as directed on the package
Cough	Robitussin cough Throat lozenges Zinc lozenges	Use as directed on the package
Diarrhea	Imodium	Use as directed on the package
Headache/Fever/Pain	Acetaminophen/Tylenol	Use as directed on the package 2 tablets every 4-6hr as needed
Hemorrhoids	Anusol-HC Preparation H	Use as directed on the package
Heartburn	Antacids: Maalox Mylanta Tums (EX) Rolaids 500's Pepcid Prilosec Tagamet	Use as directed on the package Take antacids ½ hour before meals and before bed
Nausea and Vomiting in Pregnancy	Vitamin B6 Unisom (active ingredient: doxylamine)	B6: 25-100mg twice a day Unisom: once nightly before bed
Motion Sickness	Dramamine	Use as directed on the package
Rash	Benadryl lotion Caladryl lotion Cortaid (hydrocortisone) cream	Use as directed on the package
Vaginal Yeast Infection	Gynazole 1 Gyne-Lotramin Monistat	Use as directed on the package
Sleep/Insomnia	Melatonin Unisom Benadryl	Use as directed on the package